arXiv:hep-th/9303165v1 30 Mar 1993

March, 1993

KHTP-93-02

SNUCTP-93-17

Classical $W_3^{(2)}$ algebra and its Miura map

B.K. Chung, K.G. Joo, and Soonkeon Nam*

Research Institute for Basic Sciences

and Department of Physics,

Kyung Hee University, Seoul, 130-701, Korea

Abstract

We verify that the fractional KdV equation is a bi-hamiltonian system using the zero curvature equation in $SL(3)$ matrix valued Lax pair representation, and explicitly find the closed form for the hamiltonian operators of the system. The second hamiltonian operator is the classical version of the $W_3^{(2)}$ algebra. We also construct systematically the Miura map of $W_3^{(2)}$ algebra using a gauge transformation of the $SL(3)$ matrix valued Lax operator in a particular gauge, and construct the modified fractional KdV equation as hamiltonian system.

* E-mail address: nam@nms.kyunghee.ac.kr

In the study of integrable models[1] as bi-hamiltonian systems, one of the interesting thing is the connection between the integrable nonlinear differential equations in two dimensions and the two dimensional conformal field theories. The second hamiltonian structure of the $SL(N)$ extended KdV equation corresponds to the classical form of the W_N conformal algebra[2, 3]. The corresponding integrable hierarchy of nonlinear evolution equations is characterized by a scalar Lax operator of order N, involving fields of integer conformal dimensions ranging from 2 to N[4, 5]. By the method of hamiltonian reduction, the W_N algebra is derived from an $SL(N)$ matrix in a special gauge, in which all the elements of the strictly upper triangular part are set equal to zero except those on the first diagonal, where all elements are set equal to one[6]. The classical W_3 algebra of Zamolodchikov[7], involving fields of spin-3 local current, can be obtained in the $SL(3)$ case and the corresponding integrable nonlinear differential equation is the Boussinesq(Bsq) equation.

In studying two dimensional quantum gravity, Polyakov[8] has noted that one can choose a different gauge in the $SL(3)$ case, in which the first diagonal elements of the strictly upper triangular part are set equal to zero and the second diagonal element is set equal to one. Here, in the so called $W_3^{(2)}$ algebra, two bosonic fields of spin-3/2 and a spin-1 field appear, in addition to the energy momentum tensor. This suggests that there is a whole new class of W_N algebras denoted as $W_N^{(l)}$, obtained by setting the elements of the l-th upper diagonal of an $SL(N)$ matrix to one and setting the other elements in the upper triangular part to zero.

As the W_N algebra are directly related to the second hamiltonian structure of $SL(N)$ extended KdV equation, it is a natural that this will also be the case for the $W_N^{(l)}$ algebras[9, 11]. It was shown in Ref.[9] that the new non-linear evolution equations, called the fractional KdV equation, can be constructed by using dimensional reduction of self-dual Yang-Mills equations with two killing symmetries, and making suitable ansätz for the matrix components of the gauge connections. These equation can be written as hamiltonian systems with poisson brakets given by the classical $W_3^{(2)}$ algebra. These equations were found to be different from the Bsq equation, for which the

second hamiltonian structure is the standard W_3 algebra. Such an equation can be also obtained from the Bsq equation by interchanging the role of independent variables x and t[11]. Furthemore, the free field representation, i.e. the generalized Miura map, for W_N was obtained either from a gauge tranformation[6], or a factorization of the scalar Lax operator[4, 5]. Introducing a two component scalar field and two bosonic fields with spin-1/2, the free field represention for $W_3^{(2)}$ algebra was also constructed [9, 11, 12].

In this paper we will verify the fractional KdV equation as a bi-hamiltonian system using the zero curvature condition with $SL(3)$ matrix valued Lax pair representation and explicitly find the closed form for the hamiltonian operators of the system. The second hamiltonian operator is the classical version of the $W_3^{(2)}$ algebra. From this, one can formulate the non-linear equations related to $W_N^{(l)}$ conformal algebras for all the integer N and l $(1 \leq l \leq N-1)$. We will also construct systematically the Miura map of $W_3^{(2)}$ algebra using a gauge transformation of the $SL(3)$ matrix valued Lax operator and particular gauge choice, and verify the modified fractional KdV equation as hamiltonian system. This method can be extended to the general $W_N^{(l)}$ case.

First of all, we will consider the well-known hamiltonian operators and Miura map of the Bsq system in the zero curvature formulation. This will serve to fix the notations we will be using in the case of the fractional KdV system. We parametrize two gauge potentials of the $SL(3)$ matrix valued Lax pair operators as

$$U = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ u_1 + \lambda & u_2 & 0 \end{pmatrix}, \quad V = \begin{pmatrix} P_1(x,t) & P_2(x,t) & P_3(x,t) \\ Q_1(x,t) & Q_2(x,t) & Q_3(x,t) \\ S_1(x,t) & S_2(x,t) & S_3(x,t) \end{pmatrix}, \quad TrV = 0, \qquad (1)$$

where λ is a constant parameter, and the corresponding Lax operator is $L = \partial_x - U$. Using the zero curvature condition $[\partial_x - U, \partial_t - V] = 0$, we obtain the dynamical equations of u_1, u_2 and the constraint equations[13]. These equations are expressed only in terms of the variables of upper triangular part of V, i.e. P_2, P_3 and Q_3. Redefining $u = u_2, \quad v = u_{2x} - 2u_1$, we recover the standard expression[9, 11]. Using one of the constraint

equations, $P_{3x} + P_2 - Q_3 = 0$, and setting $\tilde{Q}_3 = Q_3 - \frac{1}{2}P_{3x}$, $\tilde{P}_3 = -\frac{1}{2}P_3$, we can write the hamiltonian equations of u, v variables in a matrix form:

$$\begin{pmatrix} u \\ \\ v \end{pmatrix}_t = \left(\mathcal{D}^{(2)}[u,v] - 6\lambda\mathcal{D}^{(1)}[u,v]\right) \begin{pmatrix} \tilde{Q}_3 \\ \\ \tilde{P}_3 \end{pmatrix}, \tag{2}$$

where the first hamiltonian operator, $\mathcal{D}^{(1)}[u,v]$, and the second hamiltonian operator, $\mathcal{D}^{(2)}[u,v]$, are

$$\mathcal{D}^{(1)}[u,v] = \begin{pmatrix} 0 & \partial_x \\ \\ \partial_x & 0 \end{pmatrix},$$

$$\mathcal{D}^{(2)}[u,v] = \begin{pmatrix} -2\partial_x^3 + 2u\partial_x + u_x & 3v\partial_x + 2v_x \\ \\ 3v\partial_x + v_x & \begin{array}{c} 2\partial_x^5/3 - 10u\partial_x^3/3 - 5u_x\partial_x^2 \\ +(8u^2/3 - 3u_{xx})\partial_x - 2u_{xxx}/3 + 8uu_x/3 \end{array} \end{pmatrix}. \tag{3}$$

When we put the upper triangular part of V as the dual of the upper triangular part of U, i.e. $P_2 = Q_3 = 0$, $P_3 = 1$ ($\tilde{Q}_3 = 0$, $\tilde{P}_3 = -1/2$), we obtain the Bsq equation,

$$u_t = -v_x, \quad v_t = \frac{1}{3}u_{xxx} - \frac{4}{3}uu_x.$$

The other elements of V are obtained in terms of u, v variables by the constraint equation. The hamiltonian operators of eq.(3) define the poisson structure of Bsq equation. The poisson brackets represented by the second hamiltonian operator,

$$\begin{pmatrix} \{u(x), u(y)\} & \{u(x), v(y)\} \\ \\ \{v(x), u(y)\} & \{v(x), v(y)\} \end{pmatrix} = \mathcal{D}^{(2)}[u,v]\delta(x-y), \tag{4}$$

is the classical version of W_3 algebra[3].

To look for the Miura map of the Bsq equation, that is the free field representation of the above W_3 algebra, we perform the gauge transformation $L \longrightarrow \Phi^{-1}L\Phi$ for the Lax operator L with $\Phi(x,t)$ taking values in the strictly lower triangular matrices of $SL(3)$

with the diagonal elements set equal to one. Setting the transformed Lax operator as $\tilde{L} = \partial_x - \tilde{U}$, where $\tilde{U} = \Phi^{-1}U\Phi - \Phi^{-1}\Phi_x$, and choosing a gauge as

$$\tilde{U} = \begin{pmatrix} j_1 & 1 & 0 \\ 0 & -(j_1+j_2) & 1 \\ \lambda & 0 & j_2 \end{pmatrix}, \tag{5}$$

we obtain the Miura map of u and v with respect to j_1 amd j_2,

$$\begin{aligned} u &= (j_1 - j_2)_x + j_1^2 + j_1 j_2 + j_2^2, \\ v &= -(j_1+j_2)_{xx} - (2j_1 - j_2)j_{1x} + (2j_2 - j_1)j_{2x} + 2(j_1+j_2)j_1 j_2. \end{aligned} \tag{6}$$

The modified Bsq equation(mBsq) for j_1, j_2 and its hamiltonian structure can be exhibited by the Fréchet derivative of the transformation of eq.(6). In general, given a transformation $u = F[j, j_x, \cdots]$, the Fréchet derivative, $\left[\frac{du}{dj}\right]$, is the differential operater that implies $u_t = \left[\frac{du}{dj}\right] j_t$ and for any functional $H[u]$, we have $\frac{\delta H}{\delta j} = \left[\frac{du}{dj}\right]^* \frac{\delta H}{\delta u}$, where $\left[\frac{du}{dj}\right]^*$ is the formal adjoint of $\left[\frac{du}{dj}\right]$. The Fréchet derivative and its formal adjoint with respect to the transformation eq.(6) can be written in a matrix form as

$$\begin{aligned} \left[\frac{d(u,v)}{d(j_1,j_2)}\right] &= \begin{pmatrix} \partial_x + 2j_1 + j_2 & -\partial_x + 2j_2 + j_1 \\ -\partial_x^2 + (j_2 - 2j_1)\partial_x - 2j_{1x} & -\partial_x^2 + (2j_2 - j_1)\partial_x + j_{1x} \\ -j_{2x} + 2j_2^2 + 4j_1 j_2 & +2j_{2x} + 4j_1 j_2 + 2j_1^2 \end{pmatrix}, \\ \left[\frac{d(u,v)}{d(j_1,j_2)}\right]^\dagger &= \begin{pmatrix} -\partial_x + 2j_1 + j_2 & -\partial_x^2 - (j_2 - 2j_1)\partial_x - j_{2x} + 2j_2^2 + 4j_1 j_2 \\ \partial_x + 2j_2 + j_1 & -\partial_x^2 - (2j_2 - j_1)\partial_x + 2j_{1x} + 4j_1 j_2 + 2j_1^2 \end{pmatrix}. \end{aligned} \tag{7}$$

The hamiltonian operator of mBsq equation,$\mathcal{D}[j_1, j_2]$, and the second hamiltonian operator of Bsq equation, $\mathcal{D}^{(2)}[u, v]$, are related by, through the Fréchet derivative and its formal adjoint,

$$\mathcal{D}^{(2)}[u,v] = \left[\frac{d(u,v)}{d(j_1,j_2)}\right] \mathcal{D}[j_1,j_2] \left[\frac{d(u,v)}{d(j_1,j_2)}\right]^\dagger, \tag{8}$$

and it is straightforward to show that $\mathcal{D}[j_1, j_2]$ takes the form

$$\mathcal{D}[j_1, j_2] = \frac{1}{3}\begin{pmatrix} 2\partial_x & -\partial_x \\ -\partial_x & 2\partial_x \end{pmatrix}. \tag{9}$$

(This operator can be also found introducing the inverse Fréchet derivative[14].) This hamiltonian operator defines the poisson structure of mBsq equation as eq.(4) in the case of Bsq equation. The mBsq equation for j_1,j_2 is

$$\begin{pmatrix} j_1 \\ j_2 \end{pmatrix}_t = \mathcal{D}[j_1, j_2]\left[\frac{d(u,v)}{d(j_1,j_2)}\right]^{\dagger}\begin{pmatrix} \frac{\delta}{\delta u} \\ \frac{\delta}{\delta v} \end{pmatrix} H = \frac{1}{3}\begin{pmatrix} (j_{1x} + 2j_{2x} + j_1^2 - 2j_2^2 - 2j_1 j_2)_x \\ (-2j_{1x} - j_{2x} - 2j_1^2 + j_2^2 - 2j_1 j_2)_x \end{pmatrix}, \tag{10}$$

where we have used $\frac{\delta H}{\delta u} = \tilde{Q}_3 = 0$, $\frac{\delta H}{\delta v} = \tilde{P}_3 = -\frac{1}{2}$ in eq.(2).

Through above methods, we can new find the fractional KdV equation as a bi-hamiltonian system and show that the second hamiltonian operator is the classical version of $W_3^{(2)}$ algebra, and we also find explicitly the Miura map. Firstly, we parametrize two gauge potentials of $SL(3)$ matrix valued Lax pair operators as follows:

$$U = \begin{pmatrix} \frac{1}{2}H(x,t) & 0 & 1 \\ G^+(x,t) & -H(x,t) & 0 \\ D(x,t) + \lambda & G^-(x,t) & \frac{1}{2}H(x,t) \end{pmatrix}, \quad V = \begin{pmatrix} P_1(x,t) & P_2(x,t) & P_3(x,t) \\ Q_1(x,t) & Q_2(x,t) & Q_3(x,t) \\ S_1(x,t) & S_2(x,t) & S_3(x,t) \end{pmatrix}, \tag{11}$$

with $TrV = 0$, where λ is a constant parameter. The zero curvature condition, $[\partial_t - U, \partial_x - V] = 0$, gives five constraint equations for S_1, S_2, S_3, Q_1 and Q_2, and four dynamical equations for H, G^+, G^- and D. If we redefine $D + \frac{3}{4}H^2 = T$, the four dynamical equations for the variables H, G^+, G^- and T are expressed in terms of P_1, P_2, P_3 and Q_3 after using the constraint equation as

$$H_t = 2P_{1x} + G^+P_2 - G^-Q_3 + P_{3xx} \ ,$$

$$
\begin{aligned}
G^+_t =& -3G^+P_1 - Q_{3xx} - 3HQ_{3x} + (\lambda + T - \frac{3}{2}H_x - 3H^2)Q_3 + G^+_x P_3 + \frac{3}{2}G^+ H P_3 \ , \\
G^-_t =& 3G^-P_1 + P_{2xx} - 3HP_{2x} + (-\lambda - \frac{3}{2}H_x + 3H^2 - T)P_2 + G^-_x P_3 \\
& +3G^-P_{3x} - \frac{3}{2}HG^-P_3, \\
T_t =& 3HP_{1x} + \frac{3}{2}G^+P_{2x} + \frac{1}{2}G^+_x P_2 + \frac{1}{2}G^-_x Q_3 + \frac{3}{2}G^-Q_{3x} - \frac{1}{2}P_{3xxx} \\
& +\frac{3}{2}HP_{3xx} + 2(\lambda + T - \frac{3}{4}H^2)P_{3x} + (T_x - \frac{3}{2}HH_x)P_3 \ .
\end{aligned}
\tag{12}
$$

Setting $\tilde{P}_1 = 3P_1 + \frac{3}{2}P_{3x} - \frac{3}{2}HP_3$, we can show that above eqution express the hamiltonian equations for the variables H, $G^{(\pm)}$ and T in a closed form. Splitting eq.(12) into two parts, one that depends on the constant parameter λ and the rest, one can define two hamiltonian operators, $\mathcal{D}^{(1)}$ and $\mathcal{D}^{(2)}$. That is, the eq.(12) reads in a compact form as follows:

$$
\Psi^\dagger_t = \left(\mathcal{D}^{(2)}[H, G^+, G^-, T] + \lambda \mathcal{D}^{(1)}[H, G^+, G^-, T]\right)\mathcal{P}^\dagger \, , \tag{13}
$$

where $\Psi = (H, G^+, G^-, T)$,$\mathcal{P}$=$(\tilde{P}_1, P_2, Q_3, P_3)$, and the subscript t denotes the time derivative. Here the matrix $\mathcal{P}$ means the functional variation for a hamiltonian,$\mathcal{H}$, of the equation, i.e. $\tilde{P}_1 = \dfrac{\delta}{\delta H}\mathcal{H}$, $P_2 = \dfrac{\delta}{\delta G^+}\mathcal{H}$, $Q_3 = \dfrac{\delta}{\delta G^-}\mathcal{H}$ and $P_3 = \dfrac{\delta}{\delta T}\mathcal{H}$. The first hamiltonian operator $\mathcal{D}^{(1)}$ has non-vanishing elements ${\mathcal{D}^{(1)}}_{23} = 1$, ${\mathcal{D}^{(1)}}_{32} = -1$, and ${\mathcal{D}^{(1)}}_{44} = 2\partial_x$, and the second hamiltonian operator $\mathcal{D}^{(2)}$ is

$$
\mathcal{D}^{(2)} = \begin{pmatrix}
\frac{2}{3}\partial_x & G^+ & -G^- & H_x + H\partial_x \\
-G^+ & 0 & \begin{matrix}-\partial_x^2 - 3H\partial_x + T \\ -3H_x/2 - 3H^2\end{matrix} & \frac{3}{2}G^+\partial_x + G^+_x \\
G^- & \begin{matrix}\partial_x^2 - 3H\partial_x - T \\ -3H_x/2 + 3H^2\end{matrix} & 0 & \frac{3}{2}G^-\partial_x + G^-_x \\
H\partial_x & \frac{3}{2}G^+\partial_x + \frac{1}{2}G^+_x & \frac{1}{2}G^-_x + \frac{3}{2}G^-\partial_x & -\frac{1}{2}\partial_x^3 + 2T\partial_x + T_x
\end{pmatrix} . \tag{14}
$$

Putting $\mathcal{P}$=(0,1,1,0) in eq.(13), the other elements of V are fixed by the constraint equations as $P_1 = Q_2 = S_3 = 0$, $Q_1 = S_2 = -\frac{3}{2}H$ and $S_1 = \frac{1}{2}(G^+ + G^-)$, and we can recover the fractional KdV equation of Ref.[9], which we will call the 'first' hamiltonian equation, and see that the hamiltonian of the equation is $\mathcal{H} = \int (G^+ + G^-)\, dx$. In this

particular case, the upper triangular part of V is a dual of the upper triangular part of U. Returning to the general form of $\mathcal{D}^{(1)}$ and $\mathcal{D}^{(2)}$, the non-vanishing poisson brackets of the first hamiltonian structure can be written as

$$\{G^+(x), G^-(y)\} = \delta(x-y), \quad \{T(x), T(y)\} = 2\partial_x\delta(x-y),$$

and for the second structure they are as follows:

$$\begin{aligned}
\{H(x), H(y)\} &= \frac{2}{3}\delta(x-y), \quad \{T(x), H(y)\} = H(x)\partial_x\delta(x-y)\ , \\
\{H(x), G^{(\pm)}(y)\} &= \pm G^{(\pm)}(x)\delta(x-y), \quad \{G^{(\pm)}(x), G^{(\pm)}(y)\} = 0\ , \\
\{T(x), G^{(\pm)}(y)\} &= (\frac{3}{2}G^{(\pm)}(x) + \frac{1}{2}G^{(\pm)}(x))\delta(x-y), \\
\{T(x), T(y)\} &= (-\frac{1}{2}\partial_x^3 + 2T(x)\partial_x + T_x(x))\delta(x-y)\ , \\
\{G^+(x), G^-(y)\} &= (-\partial_x^2 - 3H(x)\partial_x + T(x) - \frac{3}{2}H_x(x) - 3H^2(x))\delta(x-y)\ .
\end{aligned} \tag{15}$$

These poisson brackets are the classical version of $W_3^{(2)}$ algebra involving an abelian spin-1 current H, a spin-2 field T (stress-energy tensor) and two bosonic spin-3/2 fields $G^{(\pm)}$, which were originally introduced by Polyakov[8] and further investigated by Bershadsky[12].

It is important that find the Miura map of hamiltonian equation(13), i.e. the free field representation, if we want to quantize the hamiltonian system. We now look for the Miura map of fractional KdV equation systematically and show that it is the free field realization of the above second poisson brackets, also find the modified fractional KdV equations as hamiltonian system. As in the case of W_3, we perform the gauge transformation for the Lax operator L with $\Phi(x,t)$ taking values in the strictly lower triangular matrices of $SL(3)$ with ones on the diagonal. The gauge potential $\tilde{U}$ of the transformed Lax operator $\tilde{L}$ is $\tilde{U} = \Phi^{-1}U\Phi - \Phi^{-1}\Phi_x$, and when we choose a gauge as

$$\tilde{U} = \begin{pmatrix} j_3 & j_1 & 1 \\ 0 & -(j_3+j_4) & j_2 \\ \lambda & 0 & j_4 \end{pmatrix}\ , \tag{16}$$

we obtain the Miura map;

$$H = j_1 j_2 + j_3 + j_4, \quad T = j_3^2 + j_4^2 + j_3 j_4 - \frac{1}{2} j_1 j_{2x} + \frac{1}{2} j_2 j_{1x} + \frac{1}{2} j_{3x} - \frac{1}{2} j_{4x} \ ,$$
$$G^+ = -j_{2x} - 2 j_2 j_3 - j_1 j_2^2 - j_2 j_4, \quad G^- = j_{1x} - j_1^2 j_2 - 2 j_1 j_4 - j_1 j_3 \ . \tag{17}$$

We can now see that j_1, j_2 are bosonic fields of spin half, and j_3, j_4 have spin 1. Here we can clearly see the advantage of the redefinition of $D + \frac{3}{4} H^2 = T$. In terms of j's D has quartic terms, whereas T requires only those quadratic in j's. The latter is suitable for free field representation, which is easier to quantize. In fact, to show that the above Miura map is the free field realization of $W_3^{(2)}$ algebra, we must find the poisson structure of j's.

The hamiltonian operator of the modified fractional KdV equation,$\mathcal{D}[j_1, j_2, j_3, j_4]$, which defines the poisson structure as before, and the second hamiltonian operator of the fractional KdV are related by, through the Fréchet derivative and its formal adjoint,

$$\mathcal{D}^{(2)}[H, G^+, G^-, T] = \left[\frac{d(H, G^+, G^-, T)}{d(j_1, j_2, j_3, j_4)} \right] \mathcal{D}[j_1, j_2, j_3, j_4] \left[\frac{d(H, G^+, G^-, T)}{d(j_1, j_2, j_3, j_4)} \right]^\dagger \ . \tag{18}$$

The Fréchet derivative and its formal adjoint with respect to the transformation eq.(17) are explicity calculated as :

$$\left[\frac{d(H, G^+, G^-, T)}{d(j_1, j_2, j_3, j_4)} \right] = \begin{pmatrix} j_2 & j_1 & 1 & 1 \\ -j_2^2 & \begin{matrix} -\partial_x - 2j_3 \\ -2j_1 j_2 - j_4 \end{matrix} & -2j_2 & -j_2 \\ \begin{matrix} \partial_x - 2j_1 j_2 \\ -2j_4 - j_3 \end{matrix} & -j_1^2 & -j_1 & -2j_1 \\ -\frac{1}{2} j_{2x} + \frac{1}{2} j_2 \partial_x & -\frac{1}{2} \partial_x + \frac{1}{2} j_{1x} & \begin{matrix} \partial_x/2 \\ +2j_3 + j_4 \end{matrix} & \begin{matrix} -\partial_x/2 \\ +2j_4 + j_3 \end{matrix} \end{pmatrix} ,$$

$$\left[\frac{d(H, G^+, G^-, T)}{d(j_1, j_2, j_3, j_4)} \right]^\dagger = \begin{pmatrix} j_2 & -j_2^2 & \begin{matrix} -\partial_x - 2j_1 j_2 \\ -2j_4 - j_3 \end{matrix} & j_{2x} - \frac{1}{2} j_2 \partial_x \\ j_1 & \begin{matrix} \partial_x - 2j_3 \\ -2j_1 j_2 - j_4 \end{matrix} & j_1^2 & \frac{1}{2} j_1 \partial_x + j_{1x} \\ 1 & -2j_2 & -j_1 & 2j_3 + j_4 - \frac{1}{2} \partial_x \\ 1 & -j_2 & -2j_1 & 2j_4 + j_3 + \frac{1}{2} \partial_x \end{pmatrix} . \tag{19}$$

Using eq.(18) and the second hamiltonian operator of eq.(14), it is straightforward to show that the hamiltonian operator of modified fractional KdV equation is

$$\mathcal{D}[j_1, j_2, j_3, j_4] = \begin{pmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & \frac{2}{3}\partial_x & -\frac{1}{3}\partial_x \\ 0 & 0 & -\frac{1}{3}\partial_x & \frac{2}{3}\partial_x \end{pmatrix} . \tag{20}$$

In particular we can see that the hamiltonian operator of mBsq equation eq.(9) is exactly involved in it of modified fractional KdV equation. The above operator defines the poisson brakets of the modified fractional KdV system, of which non-vanishing ones are

$$\begin{aligned} &\{j_1(x), j_2(y)\} = \delta(x-y)\,, \quad \{j_3(x), j_3(y)\} = \frac{2}{3}\partial_x\delta(x-y)\,, \\ &\{j_3(x), j_4(y)\} = -\frac{1}{3}\partial_x\delta(x-y)\,, \quad \{j_4(x), j_4(y)\} = \frac{2}{3}\partial_x\delta(x-y)\,. \end{aligned}$$

Setting $\phi_1 = \frac{\sqrt{3}}{2}(j_3 + j_4), \phi_2 = \frac{1}{2}(j_3 - j_4)$, we can obtain the poisson brackets in terms of j_1, j_2 and ϕ_1, ϕ_2 that express the free field realization.

Finally, we find the modified fractional KdV equation as hamiltonian system using the hamiltonian operator eq.(20) and the Fréchet derivative eq.(19). The hamiltonian equation for j's are

$$J_t^\dagger = \mathcal{D}[j_1, j_2, j_3, j_4]\left[\frac{d(H, G^+, G^-, T)}{d(j_1, j_2, j_3, j_4)}\right]^\dagger \Delta^\dagger \mathcal{H}\,, \tag{21}$$

where $J = (j_1, j_2, j_3, j_4)$, $\Delta = (\frac{\delta}{\delta H}, \frac{\delta}{\delta G^+}, \frac{\delta}{\delta G^-}, \frac{\delta}{\delta T},)$ and $\mathcal{H}$ is a conserved quantity of the fractional KdV equation, therefore $\Delta\mathcal{H}$ is the very $\mathcal{P}$ in eq.(13). When we take $\Delta\mathcal{H}$=$\mathcal{P}$= $(0, 1, 1, 0)$ which give the fractional KdV equation of Ref.[9], we can obtain a modified fractional KdV equation following as

$$\begin{aligned} &j_{1t} = -j_1^2 - 2j_3 - 2j_1j_2 - j_4\,, \quad j_{2t} = j_2^2 + 2j_1j_2 + j_3 + 2j_4\,, \\ &j_{3t} = -2j_{2x}, \quad j_{4t} = -2j_{1x}\,. \end{aligned} \tag{22}$$

Using the above-mentioned variables ϕ_1, ϕ_2 and resetting $p_1 = \frac{1}{\sqrt{2}}(j_1 - j_2)$, $p_2 = \frac{1}{\sqrt{2}}(j_1 + j_2)$, these equations agree with the equation (called $\widetilde{mBsq}$) given in Ref.[11] except the factors.

As a final remark, we will discuss a systematic description of the $W_N^{(l)}$ algebras and its Miura map for all N and l ($1 \leq l \leq N-1$). For this we have to parametrize the gauge potential defining the $SL(N)$ matrix valued Lax operator U, similar to U of eq.(1) and eq.(11) for the case of $SL(3)$, where the elements of the l-th upper diagonal part are set equal to one and the other elements in the upper triangular part to zero[11]. However this parametrization for $N > 3$ have not been explicitly constructed yet. The zero curvature condition expresses dynamical equations for the elements of U and constraint equations for the elements of V. Using the constraint equations through an appropriate change of variables in V, the dynamical equations will give hamiltonian equations, and the hamiltonian operator will represent the classical $W_N^{(l)}$ algebra. As a particular case, when we take the upper triangular part of V same as the upper triangular part of U, i.e. if we make use of the $x \leftrightarrow t$ duality, the hamiltonian equation will represent the 'first' non-linear differential equation of corresponding algebra. The Miura map of this non-linear differential equation represents the free field realization of corresponding algebra. It is well known that the Miure map for W_N algebra ($l = 1$ in $W_N^{(l)}$ case) is nothing but the relation between two different choice of gauge slice[4, 6]. In this paper we also found the Miura map for $W_3^{(2)}$ case using an appropriate gauge choice. Thus, we expect such a procedure will also gives Miura maps of general $W_N^{(l)}$.

We would like to thank Q-Han Park for discussions. This work was supported in part by Ministry of Education, and by Korea Science and Engineering Foundation.

References

[1] L.D. Fadeev and L.A. Takhtajan, Hamiltonian Methods in the Theory of Solitons (Springer, Berlin, 1987); A. Das, Integrable Models (World Scientific, Singapore, 1989); V.E. Zakharov, S.V. Manakov, S.P. Novikov and L.P. Pitaievski, Theory of Solitons.(New York, Plenum 1984).

[2] J.L. Gervais, Phys. Lett. **B160** (1985) 277, 279.

[3] P. Mathieu, Phys. Lett. **B208** (1988) 101.

[4] V.G. Drinfeld and V.V. Sokolov, J. Sov. Math. **30** (1985) 1975.

[5] B.A. Kupershmidt and G. Wilson, Inv. Math. **62** (1981) 403.

[6] M. Bershadsky and H. Ooguri, Commun. Math. Phys. **126** (1989) 49.

[7] A.B. Zamolodchikov, Theor. Math.Phys. **654** (1985) 1205.

[8] A.M. Polyakov, Int. J. Mod. Phys. **A5** (1990) 833.

[9] I. Bakas and D. Depireux. Mod. Phys. Lett. **A6** (1991) 399; Int. J. Mod. Phys. **A7** (1992) 1767; Mod. Phys. Lett. **A6** (1991) 1561 (erratum *ibid.* **A6** (1991) 2351).

[10] N.J. Burroughs, M.F de Groot, T.J. Hollowood and J.L. Miramontes. "Generalized Drinfel'd-Sokolov hierarchies II : The hamiltonian structures" Princeton preprint, PUPT-1263, IASSNS-HEP-91/42 (1991).

[11] P. Mathieu and W. Oevel, Mod. Phys. Lett. **A6** (1991) 2397.

[12] M. Bershadsky, Commun. Math. Phys. **139** (1991) 71.

[13] A. Das, W.J. Huang, and S. Roy, Phys. Lett. **A153** (1991) 186.

[14] B.K. Chung, K.G. Joo, and Sookeon Nam, Kyung Hee preprint, KHTP-93-03 (March, 1993).